



Multivariate approach in *Eucalyptus* and its effects in the genotypes × environments interaction

Osmarino Pires Dos Santos¹, Ivan Ricardo Carvalho², Vinícius Jardel Szareski², Alan Junior De Pelegrin¹, Mauricio Horbach Barbosa¹, Francine Lautenchleger³, Vélci, Queiróz De Souza⁴

¹Centro de Genômica e Fitomelhoramento, Universidade Federal de Pelotas, Capão do Leão, RS, Brazil

²Departamento de Fitotecnia, Universidade Federal de Pelotas, Capão do Leão, RS, Brazil

³Departamento Agronomia, Universidade Estadual de Maringá, Maringá, PR, Brazil.

⁴Universidade Federal do Pampa, Dom Pedrito, RS, Brazil.

Corresponding author: Ivan Ricardo Carvalho

E-mail: carvalho.irc@gmail.com

Genet. Mol. Res. 17 (2): gmr16039919

Received Mar 06, 2018

Accepted May 10, 2018

Published May 15, 2018

DOI: <http://dx.doi.org/10.4238/gmr16039919>

Copyright © 2018 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution ShareAlike (CC BY-SA) 4.0 License.

ABSTRACT. The study aimed at the selection and evaluation of *Eucalyptus* spp. clones for the production of wood by means of multivariate statistical techniques. A total of 84 *Eucalyptus* clones were used belonging to the breeding program in Rio Grande do Sul, Brazil, in a randomized block design, with 20 replicates. The selection of clones from their values of genetic divergence and heterogeneity in relation to the others can be used to increase variability in breeding populations. Improvement strategies such as the Intrapopulation Recurrent Selection can be "fed" from selections made based on these parameters, but also focusing on increasing productivity and improving wood technology. There is a great difference in behavior of the clones for each environment, demonstrating the influence of the genotype × environment interaction, which can be evaluated later in the clonal tests amplified or in the observation plantings of these clones that will serve for the future indication of genotypes for the commercial plantations

Key words: *Eucalyptus* breeding; Variability genetic; Differential effects of the variation factors.

INTRODUCTION

The genus *Eucalyptus* belongs to the family Myrtaceae, originating in Australia, where it comprises more than 600 different species. In Brazil, *Eucalyptus* plantations occupy 5.7 million hectares of planted trees and are mainly located in Minas Gerais (24%), São Paulo (17%) and MatoGrosso do Sul (15%). There was a growth of 2.4% in the area of *Eucalyptus* planted in the last 5 years in the country. The state of MatoGrosso do Sul led this growth, registering an increase of 400 thousand hectares in this period, with an average growth rate

of 13% per year (Ibá, 2017). *Eucalyptus* is the main responsible for placing the country among the world's largest producers of cellulose. The increase of *Eucalyptus* productivity is mainly due to the performance of genetic improvement, through the acquisition of intra and interspecific hybrids, naturally or controlled through cloning (Fonseca et al., 2010; Rosado et al., 2012). The clonal *Eucalyptus* planting areas have been expanded throughout the national territory, due to the ease of the multiplication processes, the possibility of solving the heterogeneity of the forest plantations, as well as the promotion of the homogeneity of the technological properties of the wood, reflecting in the industrial performance improvement. Before the final recommendation of a clone, it undergoes a series of evaluations in different growing environments (Sudaricet al., 2005). This is necessary, due to the environmental diversity that occurs, most often among different growing environments, favoring the phenotypic variation of the genotypes, through the interaction between genotype and environment (GxE) (Resende, 2005). The choice of genotypes with high productivity, adaptability and stability is an important alternative to minimize the effects of the GxE interaction (Cruz & Carneiro, 2003).

In order to evaluate the physical, chemical, mechanical and anatomical properties of wood, the multivariate analyzes are widely used in structural analyzes of forests in phytosociological studies (Assiset al., 2012; Lima et al., 2011; Lobão et al., 2011; Protásio et al., 2012; Trugilho et al., 2003). Among the multivariate techniques most used by researchers are the analysis of canonical variables, grouping methods and main components. The analysis of canonical variables is one of the most suitable multivariate techniques in situations in that many dependent and independent characters of one or more genotypes are analyzed, allowing the analysis of the structure of linear relationships between two groups or character sets formed by metric variables, in order to maximize the correlation between them (Protásio et al., 2012; Hair Junior et al., 2009). The optimization method proposed by Tocher, performs the stratification of the genotypes into groups at one time, using a single criterion to keep the intra-group mean distance always lower than any intergroup distance (Cruz & Carneiro, 2006). The method proposed by Sing (1981) makes it possible to quantify the relative contribution of the characters through the genetic distance among them. Dissimilarity measures have been proposed for the quantification of distances among genotypes. The generalized distance of Mahalanobis is widely used when experiments with replicates are available (Simon et al., 2012). This is different from the other techniques because it takes into account the correlations among the characters evaluated (Cruz; Regazzi; Carneiro, 2012). In view of the need to obtain genetically superior *Eucalyptus* clones that promote productivity coupled with quality, the present study aimed at the selection and evaluation of *Eucalyptus* spp. clones for the production of wood by means of multivariate statistical techniques.

MATERIAL AND METHODS

Growing environments

The experiments were installed in three environments, in the districts of Eldorado do Sul, Butiá and São Gabriel (Table 1). As a common control in all trials, a clone of the commercial species *Eucalyptus saligna* was used.

Table 1. Geographic location and edaphoclimatic conditions of the Eldorado do Sul - RS, Butiá - RS and São Gabriel - RS environments

Municipalities	Eldorado do Sul - RS	Butiá - RS	São Gabriel - RS
Geographical coordinates	30° 06' 89" S e 51° 44' 70" W	30° 27' 50" S e 52° 10' 05" W	30° 44' 70" S e 54° 53' 42" W
Altitude (meters)	75	188	139
Previous occupation	Forest	Forest	Native field
Soil type	Gleysolic dystrophic yellow red argisol	Typic dystrophic yellow red argisol	Typical dystrophic grayish argisol
Average temperature (°C)	18.2	18.2	20.2
Frost risk	Low	Low	High
Relative humidity (%)	82.2	82.2	72.4
Annual rainfall (mm)	1570	1570	1965

Experimental conditions and genotypes used

A total of 84 *Eucalyptus* clones were used (Table 2) belonging to the breeding program in Rio Grande do Sul, Brazil, in a randomized block design, with 20 replicates and one plant per plot, spacing 3.50 × 2.14 m. Silvicultural practices were performed according to the particularities required by each environment.

The clones were evaluated at 36 months for diameter at breast height (DBH, cm), total height (H, m), total individual volume with bark (Vol, m³) and survival of clones. The total individual volume with bark was calculated using the expression: $Vol = 0.004761 + 0.000033 \times DBH^2 \times H^2$ (Table 2).

Table 2. Description of the 84 clones evaluated in the three environments and their respective species/crosses.

Number	Species/Hybrid	Number of Clones	% Of clones by species and hybrid
1	<i>E. (grandis × urophylla) × not informed</i>	31	36.90%
2	<i>E. urophylla × E. globulus</i>	21	25.00%
3	<i>E. (grandis × urophylla) × E. globulus</i>	12	14.29%
4	<i>E. saligna × not informed</i>	4	4.76%
5	<i>E. (grandis × urophylla) × E. (urophylla × globulus)</i>	3	3.57%
6	<i>E. (grandis × saligna) × not informed</i>	2	2.38%
7	<i>E. grandis × E. globulus</i>	2	2.38%
8	<i>E. (grandis × urophylla) × E. maidenii</i>	2	2.38%
9	<i>E. (grandis × urophylla) × E. viminalis</i>	2	2.38%
10	<i>E. urophylla × not informed</i>	1	1.19%
11	<i>E. (dunnii × grandis) × E. (urophylla × globulus)</i>	1	1.19%
12	<i>E. (dunnii × grandis) × E. viminalis</i>	1	1.19%
13	<i>E. saligna</i>	1	1.19%
14	<i>E. (dunnii × grandis) × not informed</i>	1	1.19%
Total		84	100.00%

Statistical analysis

After obtaining the data, they were submitted to a joint analysis of variance, and the significance was interpreted by the F test at 5% of probability. Afterwards, the Shapiro-Wilk test for normality proposed in 1965, based on the W statistic was performed. In order to verify the homoscedasticity of the data, the Hartley test was applied. Interaction tests were performed and then multivariate models were used. All statistical procedures were performed using the Genes software (Cruz, 2013).

RESULTS AND DISCUSSION

The analysis of variance revealed interaction, environment × clone, for all variables, diameter at breast height (DBH), plant height (PH) and volume of wood (VOM) (Table 3).

The contribution of the interaction for the phenotypic manifestation, for the variable diameter at breast height, was 21.3%; among the variation factors, the environmental parameter contributed 13.5%, and clone with 34.5%. For wood volume, the contribution of the interaction was of 18.1%, the environment 24.4% and clone with 28.4%. For these two variables the greatest contributions are in the clone variation source, demonstrating the relevance of this parameter for these variables. For the plant height variable the greatest contribution was expressed in the environment parameter, 35.2%. The interaction contributed with 15.4% and the clone variation factor, 14.9% (Table 3).

Table 3. Summary of analysis of variance for diameter at breast height (DBH), plant height (PH) and wood volume (VOM)

VF	DF	Mean Square		
		DBH	PH	VOM
Environment (A)	2	3864,89*	10261,66*	2,79*
Clone (B)	83	224,90*	93,36*	0,07*
A × B	166	74,01*	49,43*	0,02*
Block (A)	57	9,80*	12,59*	0,0039*
Residue	3552	6,49	7,01	0,0025
CV(%)		18,74	17,37	41,31
R ²		0,63	0,60	0,65

* Significant at 5% by the F-test.

Regarding the coefficient of variation, the results found (18.7%) were in agreement with Trugilho's work, where the author evaluating *Eucalyptus* clones found 21.57% for the variable diameter at breast height. Plant height (PH) showed a coefficient of 17.37%, a result superior to those described by Trugilho et al. (2001) (11.4%), working with clones, Rosado et al. (2010) (7.45%) studying *E. urophylla*, and Freitas et al. (2009) with *E. urograndis* (9.1%). The volume of wood (VOM) (41.31%) showed lower coefficient and similar to the studies conducted by Trugilho et al. (2001).

The use of multivariate analyzes aims to show the similarity among the treatments, through the use of the association among the variables, in order to form groups and to demonstrate the contribution of each character to the differentiation of the genotypes. Dismemberment of the multivariate analysis by environment allows to reveal the behavior of the clones in each environment separately.

From Table 4, it is possible to identify that, for the variable diameter at breast height (DBH), both Eldorado do Sul and São Gabriel locations obtained seven classes, where the 15 cm measurement was the most frequent. These values indicate the good performance of the clones, since more than 50% of the clones obtained values equal or superior to 15 cm. On the other hand, the Butiá environment presented a greater number of classes (eight), obtaining a lower magnitude of the values for this variable, where 12.75 cm was the most frequent measure. Also, for the variable plant height (PH), the environments differed by the number of classes, where Eldorado do Sul presented eight classes, Butiá seven classes and São Gabriel nine classes, and also by the magnitude of the classes, where the most frequent measurements in the environments of Eldorado do Sul, Butiá and São Gabriel were 18.75 cm, 13 cm, 16.5 cm respectively.

The distribution of frequencies of the phenotypic classes, shown in Figure 1, reveals that Butiá was the environment that contributed the most to the dissimilarity of the variables diameter at breast height and plant height, with 77.79% and 9.49%, respectively. For the wood volume variable, the Eldorado do Sul environment had the largest contribution to distinguish it with 39.54%, followed by the São Gabriel environment with 28.93%. (Figures 1 and 2).




Figure 1. Frequency distribution of phenotype classes for diameter at breast height (DBH), plant height (PH) and wood volume (VOM) in São Gabriel - RS, Eldorado do Sul - RS and Butiá - RS environments.




Figure 2. Frequency distribution of phenotype classes for diameter at breast height (DBH), plant height (PH) and wood volume (VOM).

According to Table 4, it is possible to verify the groupings of the genotypes by the Tocher optimization method, where, for the Eldorado do Sul environment, 13 groups were formed. It was verified that group I was composed of 27 accessions, with approximately 32% of individuals, group II with 19 accessions (22%) and group III with 16 accessions (19%), totaling 73% of the individuals, these values indicate that there is great intragroup homogeneity. Also, 27% of the remaining individuals were distributed in 10 groups, where group IV was formed by six accessions, groups VI and VII constituted by three accessions, and groups V, VIII and IX composed by two accessions. It is possible to verify that the groups X, XI, XII, XIII were composed of only one accession each, which, according to Vasconcelos et al. (2007) indicates high divergence among clones for this environment. This divergence is of great importance for the work of the breeder, since it allows its exploration for the future development of cultivars (Table 4).

Table 4. Results for the grouping of 84 *Eucalyptus* genotypes in three environments, using the Tocher optimization method, based on standardized mean Euclidean distance.

Eldorado do Sul – RS	
Groups	<i>Eucalyptus</i> accessions
I	26 53 42 33 69 52 36 54 82 39 76 79 67 60 4 18 8 40 29 51 74 84 73 32 34 12 77
II	13 24 46 71 83 70 10 59 63 16 30 62 35 75 15 65 58 11 1
III	49 50 80 57 28 23 56 20 43 78 61 27 48 21 7 3
IV	22 31 37 14 55 47
V	17 41
VI	2 68 66
VII	25 44 38
VIII	6 19
IX	9 72
X	64
XI	81
XII	5
XIII	45
Butiá –RS	
Group	<i>Eucalyptus</i> accessions
I	11 56 26 31 77 72 83 2 23 25 12 34 67 27 69 4 50 44 74 7 17 58 80 6 60 63 16 9 19 13 32 79 29 70 40 78 49 82 68 20 1 14 75
II	36 53 35 37 18 5 47 39 48 43 21
III	59 66 8 62 55
IV	45 54 28 61 30 57 41 3 46 73 81
V	33 65 71 52 64
VI	22 42
VII	10 76
VIII	51
IX	15
X	84
XI	38
XII	24
São Gabriel – RS	

Group	<i>Eucalyptus</i> accessions
I	13 79 8 24 4 30 29 18 40 47 17 51 54 16 60 26 12 69 53 2 21 52 1 84 67 71 46 76 82 5 80 58 78 61 33 39 20 35 56 31 42 27 43 50 73 7 57 3 22 74 75 62 83 10 14 32 70 41 28 77 59 34 9 63 36 55 65 68 11 72 48 19 6 15 37 64 81 49 25 38 44 45
II	23
III	66

The Butiá environment revealed grouping results similar to Eldorado do Sul, presenting the formation of 12 groups. Among the groupings, the group I highlights, composed of 43 accessions with 51.2% of the individuals. Groups II and IV were composed of 11 accessions each, groups III and V by five accessions each, groups VI and VII by two accessions each. The remaining groups (VIII, IX, X, XI and XII) were composed of clones with higher dissimilarities, where each accession formed a specific group. The results found for these two environments coincide with a study by Caixeta et al. (2003) in *Eucalyptus* clones, where the authors analyzed 44 clones, which were grouped in 11 different groups. On the other hand, the São Gabriel environment had less influence on the dissimilarity of the clones, having grouped them into only three groups, and group I was composed of 82 accessions or 97.6% of the individuals. The other two groups had only one accession each.

Through the dendrograms it is possible to perceive the genetic divergence among the 84 *Eucalyptus* clones, where the most dissimilar clones for the Eldorado do Sul environment were genotypes 26 and 45 (Figure 3). In the Butiá environment, the largest distances were found for clones 11 and 76 (Figure 4) and, for the São Gabriel environment, clones 13 and 66 (Figure 5).




Figure 3. Dendrogram with genetic divergence in 84 *Eucalyptus* genotypes in Environment 1, using normalized Euclidean distance, obtained by the nearest neighbor method.




Figure 4. Dendrogram with genetic divergence in 84 *Eucalyptus* genotypes in Environment 2, using normalized Euclidean distance, obtained by the nearest neighbor method.




Figure 5. Dendrogram with genetic divergence in 84 *Eucalyptus* genotypes in Environment 3, using normalized Euclidean distance, obtained by the nearest neighbor method.

It is possible to highlight the divergence of the environments studied from the results shown in Figures 5 and 6, which demonstrate the behavior of the clones in each growing environment. Clones that disaggregate from the others indicate greater genetic variability, thus, for the Eldorado do Sul environment, 4 groups can be distinguished, clones being clustered distinctly: clones 45, 44 and 25, clones 38, 6 and 19, clones 72 and 9, and the remainder of the clones clustered. Also, for the Butiá environment, it is possible to highlight the similar behavior of clones 22 and 24, these forming a group. Two other groups can be identified, one of which is formed by clones 76 and 10, and the other by clones 55 and 16. The São Gabriel environment had clone 66 demonstrating great variability, forming a group. Two other groups varied from the others, the first one being formed by clones 23 and 45 and the second by clones 25 and 81. Clones highlighted in distinct groups are the ones that most contribute to the interaction between genotype and environment (GXE).

Figure 5 represents the graphical dispersion in the two-dimensional space of the 85 clones under study, using the first and second canonical variables. The groups formed by means of the graphic dispersion of the scales show that in the São Gabriel environment, clone 66 was the most dissimilar among the clones tested in this environment, following the same tendency of the groups found by the Tocher optimization method and normalized Euclidean distance, thus reinforcing the data obtained by the different multivariate techniques for this environment. However, for the other environments the results obtained by the different tests were not concordant, where for the Eldorado do Sul environment, from the graphic dispersion of canonical variables it was possible to identify a more dissimilar group of clones 6, 19, 25, 38, 44 and 48, and for the Butiá environment a group with clones 76 and 10 (Figure 6).




Figure 6. Dispersion graph of 85 *Eucalyptus* clones in relation to the scores of the first two canonical variables (VC1 and VC2) for the environments of Eldorado do Sul - RS (A), Butiá (B) and São Gabriel (C).

CONCLUSION

The selection of clones from their values of genetic divergence and heterogeneity in relation to the others can be used to increase variability in breeding populations. Improvement strategies such as the Intrapopulation Recurrent Selection can be "fed" from selections made based on these parameters, but also focusing on increasing productivity and improving wood technology. There is a great difference in behavior of the clones for each environment, demonstrating the influence of the genotype \times environment interaction, which can be evaluated later in the clonal tests amplified or in the observation plantings of these clones, that will serve for the future indication of genotypes for the commercial plantations.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Indústria Brasileira De Árvores. Anuário IBÁ (2017). Brasília.

Assis MR, Protásio TP, Assis CO, Trugilho PF, et al., (2012). Qualidade e rendimentos do carvão vegetal de um clone híbrido de *Eucalyptusgrandis* \times *Eucalyptusurophylla*. *Pesq. Agrop. Bras.*, Colombo, 32(71): 291-302. <https://doi.org/10.4336/2012.pfb.32.71.291>

Caixeta RP. et al. (2003).propriedades e classificação da madeira aplicadas à seleção de genótipos de *Eucalyptus*. *Revista Árvore*, Viçosa, 27(1):43-51. <https://doi.org/10.1590/s0100-67622003000100006>

Cruz CD, Carneiro PCS (2003). Modelos biométricos aplicados ao melhoramento genético. Viçosa, MG: Ed. UFV. 2: 585.

Cruz CD, Carneiro PCS (2006). Modelos biométricos aplicados ao melhoramento genético. (2nd edn). Viçosa: UFV. p. 585.

Fonseca SM da, Resende MDV de, Alfenas AC, Guimarães LM da S, et al., (2010). Manual prático de melhoramento genético do eucalipto. Viçosa: UFV. 200p.

Freitas RGD, Vasconcelos ESD, Cruz CD, Rosado AM, et al., (2009). Predição de ganhos genéticos em progênes de polinização aberta de *Eucalyptusurograndis* cultivadas em diferentes ambientes e submetidas a diferentes procedimentos de seleção. *Rev. Árvore* [online]. 2009, 33(2): 255-263. ISSN 1806-9088.

Hair Junior JF, Black WC, Babin BJ, Anderson RE, et al., (2009). Análise multivariada de dados. Porto Alegre: Bookman. p. 688.

Lima EA, Silva HD, Lavoranti OJ (2011). Caracterização dendroenergética de árvores de *Eucalyptusbenthamii*. *Pesq. Flor. Bras.*, Colombo, 31(65):09-17. <https://doi.org/10.4336/2011.pfb.31.65.09>

Lobão MS, Castro VR, Rangel A, Sarto C, et al., (2011). Agrupamento de espécies florestais por análises univariadas e multivariadas das características anatômica, física e química das suas madeiras. *ScientiaForestalis*, Piracicaba, 39(92):469-477.

Protásio TP, Tonoli GHD, Guimarães Júnior M, Bufalino L, et al., (2012). Correlações canônicas entre as características químicas e energéticas de resíduos lignocelulósicos. *Cerne*, Lavras, 18(3):433-439. <https://doi.org/10.1590/s0104-77602012000300010>

Resende MDV (2005). Melhoramento de essências florestais. In: BORÉM, A.(Ed.). Melhoramento de espécies cultivadas. Viçosa, MG: Ed. UFV. 2: 717-780.

Rosado AM, Rosado TB, Alves AA, Laviola BG, Bhering, et al., (2012). Seleção simultânea de clones de eucalipto de acordo com produtividade, estabilidade e adaptabilidade. *Pesq. Agrop. Bras.*, 47(7):966–973. <https://doi.org/10.1590/s0100-204x2012000700013>

Rosado AM, Rosado TB, Júnior MFR.R, Bhering LL, et al., (2010). Ganhos genéticos preditos por diferentes métodos de seleção em progênes de *Eucalyptusurophylla*. *Pesq. Agrop. Bras.*, 44(12): 1653-1659. <https://doi.org/10.1590/s0100-204x2009001200014>

Shapiro SS. E, Wilk MB (1965). An Analysis of Variance Test for Normality (Complete Samples). *BiometrikaTrust*, London, 52: 591–609. <https://doi.org/10.2307/2333709>

Sudaric A, Simic D, Vrataric M (2005). Characterization of genotype by environment interactions in soybean breeding programmes of southeast Europe. *Plant Breeding*, 125: 191-194. <https://doi.org/10.1111/j.1439-0523.2006.01185.x>

Trugilho PF, Lima JT, Mori FA (2003).Correlação canônica das características químicas e físicas da madeira de clones de *Eucalyptusgrandis* \times *Eucalyptussaligna*. *Cerne*, Lavras, 9(1): 66–80.

Multivariate approach in *Eucalyptus*

Trugilho PF, Lima Tarcísio J, Mori Akira F and Lino AL (2001). Avaliação de clones de *Eucalyptus* para produção de carvão vegetal. *Cerne*, 7(2).

Vasconcelos ES, Cruz CD, Bhering LL, Resende Junior MFR (2007). Método alternativo para análise de agrupamento. *Pesq. Agrop. Bras.*, 42(10): 1421-1428. <https://doi.org/10.1590/s0100-204x2007001000008>